arXiv:1005.4318v1 [math.FA] 24 May 2010

A SOLUTION OF VARIATIONAL INEQUALITY PROBLEM FOR A FINITE FAMILY OF NONEXPANSIVE MAPPINGS IN HILBERT SPACES

FARRUKH MUKHAMEDOV AND MANSOOR SABUROV

Abstract. In this paper we prove the strong convergence of the explicit iterative process to a common fixed point of the finite family of nonexpansive mappings defined on Hilbert space, which solves the the variational inequality on the fixed points set.

Mathematics Subject Classification: 46B20; 47H09; 47H10
Key words: Nonexpansive mappings, explicit iteration process; common fixed point; variational inequality.

1. Introduction

An iterative approximation of fixed points and zeros of nonlinear operators has been studied extensively by many authors to solve nonlinear operator equations as well as variational inequality problems (see e.g., [3, 4], [7], [9]-[15], [17]-[23]). A very important class of mappings is nonexpansive mappings. In particulary, iterative approximation of fixed points of nonexpansive mappings is an important subject in nonlinear functional analysis, and has many applications in various fields of mathematics, physics etc (see e.g., [2, 16, 28]).

Let H be a real Hilbert space with inner product $\langle \cdot, \cdot \rangle$, and induced norm $\|\cdot\|$. Let $T : H \to H$ be a mapping. Denote by $F(T)$ the set of fixed points of T, that is, $F(T) = \{x \in H : Tx = x\}$. Now let us recall some well-known definitions, in which will be used in this paper.

A mapping $f : H \to H$ is said to be *contraction* if there exists $\alpha \in [0, 1)$ such that for all $x, y \in H$,

$$\|f(x) - f(y)\| \leq \alpha \|x - y\|.$$

A mapping $T : H \to H$ is called *nonexpansive* if for all $x, y \in H$,

$$\|Tx - Ty\| \leq \|x - y\|.$$

A mapping $A : H \to H$ is called $\eta-$*strongly monotone* if for all $x, y \in H$,

$$\Big\langle x - y, Ax - Ay \Big\rangle \geq \eta \|x - y\|^2.$$

A mapping $A : H \to H$ is called $k-$*Lipschitzian* if for all $x, y \in H$,

$$\|Ax - Ay\| \leq k\|x - y\|.$$

Iterative methods for nonexpansive mappings are now also applicable in solving convex minimization problems (see, for example, [26] and references therein). Let K be a closed convex nonempty subset of H and $T : K \to K$ be a nonexpansive mapping such that $F(T) \neq \emptyset$. Given $u \in K$ and a real sequence $\{\alpha_n\}_{n=1}^{\infty}$ in the interval $(0, 1)$, starting with an arbitrary initial $x_0 \in K$, let a sequence $\{x_n\}_{n=1}^{\infty}$ be defined by

$$x_{n+1} = \alpha_n u + (1 - \alpha_n)Tx_n, \quad n \in \mathbb{N}. \tag{1.1}$$

Under appropriate conditions on the iterative parameter $\{\alpha_n\}$, it has been shown by Halpern [6], Lions [8], Wittmann [25] and Bauschke [1] that $\{x_n\}_{n=1}^{\infty}$ converges strongly to $P_{F(T)}u$, the projection of u to the fixed point set, $F(T)$ of T. This means that the limit of the sequence $\{x_n\}_{n=1}^{\infty}$ solves the following minimization problem:

$$\text{find } x^* \in F(T) \text{ such that } \|x^* - u\| = \min_{x\in F(T)} \|x - u\|.$$

A typical minimization problem is to minimize a quadratic function over the set of fixed points of nonexpansive mappings in a real Hilbert space and such problems go thus: find $x^* \in F(T)$ such that

$$\frac{1}{2}\Big\langle Ax^*, x^*\Big\rangle - \Big\langle x^*, u\Big\rangle = \min_{x\in F(T)} \left(\frac{1}{2}\Big\langle Ax, x\Big\rangle - \Big\langle x, u\Big\rangle\right) \tag{1.2}$$

where $u \in H$ is a fix point and A is a mapping on H which could be monotone, strongly monotone or even bounded linear strongly positive operator as the case may be.

Let $K_1, K_2, \cdots, K_N$ be N closed convex subsets of a real Hilbert space H having a nonempty intersection K. Suppose also that each C_i is a fixed point set of nonexpansive mappings $T_i : H \to H$, $i = \overline{1, N}$. Xu [26] proved strong convergence of the iterative algorithm

$$x_{n+1} = \alpha_n u + (I - \alpha_n A)T_n x_n, \quad n \in \mathbb{N}, \tag{1.3}$$

where, I is an identity mapping and $T_n := T_{n(modN)}$, to a unique solution of the quadratic minimization problem

$$\min_{x\in K} \left(\frac{1}{2}\Big\langle Ax, x\Big\rangle - \Big\langle x, u\Big\rangle\right),$$

where A is a bounded linear strongly positive operator on H and u is a given point in H.

Marino and Xu [10], proved that the iteration scheme given by

$$x_{n+1} = \alpha_n \gamma f(x_n) + (I - \alpha_n A)T x_n, \quad n \in \mathbb{N}, \tag{1.4}$$

converges strongly to a unique solution $x' \in F(T)$ of the variational inequality

$$\Big\langle (\gamma f - A)x', y - x'\Big\rangle \le 0, \quad \forall y \in F(T), \tag{1.5}$$

which is the optimality condition for the minimization problem

$$\min_{x\in F(T)} \left(\frac{1}{2}\Big\langle Ax, x\Big\rangle - h(x)\right),$$

where h is a potential function for γf (that is, $h'(x) = \gamma f(x)$ for all $x \in H$); provided $f : H \to H$ is a contraction, $T : H \to H$ is nonexpansive and the iterative parameter $\{\alpha_n\}$ satisfies appropriate conditions.

In [27], Yamada introduced the following hybrid iterative method

$$x_{n+1} = (I - \alpha_n \mu A)T x_n, \quad n \in \mathbb{N}, \tag{1.6}$$

where T is nonexpansive, A is $L-$Lipschitzian and strongly monotone operator with constant η and $0 < \mu < \frac{2\eta}{L^2}$. He proved that if $\{\alpha_n\}$ satisfies appropriate conditions, then $\{x_n\}_{n=1}^{\infty}$ converges strongly to a unique solution $x' \in F(T)$ of the variational inequality

$$\Big\langle Ax', y - x'\Big\rangle \ge 0, \quad \forall y \in F(T).$$

Recently, motivated by the iteration schemes (1.4) and (1.6), M. Tian [24] introduced the following iterative method:

$$x_{n+1} = \alpha_n \gamma f(x_n) + (I - \alpha_n \mu A)T x_n, \quad n \in \mathbb{N}. \tag{1.7}$$

Tian [24] proved that if $f : H \to H$ is a contraction, $A : H \to H$ is an $\eta-$strongly monotone mapping, $T : H \to H$ a nonexpansive mapping and the parameter $\{\alpha_n\}$ satisfies appropriate conditions, then the sequence $\{x_n\}_{n=1}^{\infty}$ converges strongly to a unique solution $x' \in F(T)$ of the variational inequality

$$\Big\langle (\gamma f - \mu A)x', y - x'\Big\rangle \le 0, \quad \forall y \in F(T).$$

His results generalized and improved the corresponding results of Marino and Xu [10] and Yamada [27].

In this paper, we extend Tian's results [24] to a finite family of nonexpansive mappings. More precisely, we consider the following iterative algorithm

$$x_{n+1} = \alpha_n \gamma f(x_n) + (I - \alpha_n \mu A) T_n x_n, \quad n \in \mathbb{N}, \tag{1.8}$$

where $I : H \to H$ is an identity mapping, $A : H \to H$ is a $k-$Lipschitzian $\eta-$strongly monotone mapping, $\{T_i\}_{i=1}^N : H \to H$ are nonexpansive mappings with $F := \bigcap\limits_{i=1}^N F(T_i) \neq \emptyset$, and $T_n := T_{n(modN)}$. Under appropriate conditions, we shall prove strongly convergence of the sequence $\{x_n\}_{n=1}^\infty$ to a unique solution of the variational inequality

$$\Big\langle (\gamma f - \mu A)x', y - x' \Big\rangle \le 0, \quad \forall y \in F.$$

Our results generalize the corresponding results of Marino and Xu [10], Tian [24], Xu [26], Yamada [27].

2. Preliminaries

The following lemmas play an important role in proving our main results.

Lemma 2.1. [26] *Let a_n be a sequence of nonnegative numbers satisfying the following condition*

$$a_{n+1} \le (1 - \alpha_n) a_n + \alpha_n \beta_n$$

where $\{\alpha_n\}_{n=1}^\infty$, $\{\beta_n\}_{n=1}^\infty$ are sequences of real numbers such that:

(i) $0 < \alpha_n < 1$ *and* $\lim\limits_{n\to\infty} \alpha_n = 0$;

(ii) $\sum\limits_{n=1}^\infty \alpha_n = \infty$;

(iii) $\limsup\limits_{n\to\infty} \beta_n \le 0$.

Then $\lim\limits_{n\to\infty} a_n = 0$

Lemma 2.2. [26] *In a real Hilbert space H, there holds the following inequality*

$$\|x + y\|^2 \le \|x\|^2 + 2\langle y, x + y\rangle,$$

for all $x, y \in H$.

Lemma 2.3. [5] *Let H be a Hilbert space and $T : H \to H$ be a nonexpansive mapping with $Fix(T) \neq \emptyset$. If $\{x_n\}_{n=1}^\infty$ converges weakly to x and $\{\|x_n - Tx_n\|\}_{n=1}^\infty$ converges to 0, then $Tx = x$.*

Lemma 2.4. [24] *Let H be a real Hilbert space. Let $I : H \to H$ be an identity mapping, $f : H \to H$ be a contraction mapping with $0 < \alpha < 1$, $A : H \to H$ be a $k-$Lipschitzian $\eta-$strongly monotone mapping, and $T : H \to H$ be a nonexpansive mapping with $F(T) \neq \emptyset$. Let x_t be a unique solution of the following equation*

$$x_t = t\gamma f(x_t) + (I - t\mu A) T x_t,$$

where $0 < \mu < \frac{2\eta}{k^2}$, $\tau := \mu(\eta - \frac{\mu k^2}{2})$, $0 < \gamma < \frac{\tau}{\alpha}$, and $0 < t < 1$. Then the following variational inequality

$$\Big\langle (\gamma f - \mu A)x, y - x \Big\rangle \le 0, \quad \forall y \in F(T). \tag{2.1}$$

has a unique solution x' on the set $F(T)$ and x_t converges strongly to $x^{'}$ as $t \to 0$.

3. Main results

In this section we shall prove our main results. To formulate ones, we need some auxiliary results.

Lemma 3.1. *Let H be a real Hilbert space. Let $I : H \to H$ be an identity mapping, $f : H \to H$ be a contraction mapping with $0 < \alpha < 1$, $A : H \to H$ be a $k-$Lipschitzian $\eta-$strongly monotone mapping, and $\{T_i\}_{i=1}^N : H \to H$ be nonexpansive mappings with $F(T_i) \neq \emptyset$, for all $i = \overline{1, N}$. If $0 < \mu < \frac{2\eta}{k^2}$ and $0 < \alpha_n < 1$ for all $n \in \mathbb{N}$ then we have the following inquality*

$$\|(I - \alpha_n \mu A)T_n x - (I - \alpha_n \mu A)T_n y\| \le (1 - \alpha_n \tau)\|x - y\|, \quad \forall n \in \mathbb{N}, \tag{3.1}$$

for all $x, y \in H$, where, $T_n := T_{n(modN)}$ and $\tau := \mu(\eta - \frac{\mu k^2}{2})$.

Proof. Let us denote by $S_n := (I - \alpha_n \mu A)T_n$. Then, we have for all $x, y \in H$ and $n \in \mathbb{N}$

$$\begin{aligned}
\|S_n x - S_n y\|^2 &\le \|T_n x - T_n y\|^2 - 2\alpha_n \mu \Big\langle T_n x - T_n y, AT_n x - AT_n y \Big\rangle \\
&\quad + \alpha_n^2 \mu^2 \|AT_n x - AT_n y\|^2 \\
&\le \left(1 - 2\alpha_n \mu \eta + \alpha_n^2 \mu^2 k^2\right) \|T_n x - T_n y\|^2 \\
&\le \left(1 - 2\alpha_n \mu \left(\eta - \frac{\mu k^2}{2}\right)\right) \|x - y\|^2 \\
&= (1 - 2\alpha_n \tau)\|x - y\|^2 \\
&\le (1 - \alpha_n \tau)^2 \|x - y\|^2,
\end{aligned}$$

which means

$$\|S_n x - S_n y\| \le (1 - \alpha_n \tau)\|x - y\|. \tag{3.2}$$

This completes the proof. □

Theorem 3.2. *Let H be a real Hilbert space. Let $I : H \to H$ be an identity mapping, $f : H \to H$ be a contraction mapping with $0 < \alpha < 1$, $A : H \to H$ be a $k-$Lipschitzian $\eta-$strongly monotone mapping, and $\{T_i\}_{i=1}^N : H \to H$ be nonexpansive mappings with $F := \bigcap_{i=1}^{N} F(T_i) \neq \emptyset$. Suppose that $0 < \mu < \frac{2\eta}{k^2}$, $\tau := \mu(\eta - \frac{\mu k^2}{2})$, $0 < \gamma < \frac{\tau}{\alpha}$, and the sequence $\{\alpha_n\}_{n=1}^\infty \subset (0, 1)$ satisfies the following condition:*

(i) $\lim\limits_{n\to\infty} \alpha_n = 0$;

(ii) $\sum\limits_{n=1}^{\infty} \alpha_n = \infty$;

(iii) $\lim\limits_{n\to\infty} \dfrac{\alpha_n}{\alpha_{n+N}} = 1$.

If $F = F(T_N T_{N-1} \cdots T_1) = F(T_1 T_N T_{N-1} \cdots T_2) = F(T_2 T_1 T_N T_{N-1} \cdots T_3) = \cdots = F(T_{N-1} T_{N-2} \cdots T_1 T_N)$ then the sequence $\{x_n\}_{n=1}^\infty$ which is defined by (1.8), converges strongly to a unique solution $x' \in F$ of the variational inequality

$$\Big\langle (\gamma f - \mu A)x', y - x' \Big\rangle \le 0, \quad \forall y \in F. \tag{3.3}$$

Proof. Since the mapping $T_N T_{N-1} \cdots T_1 : H \to H$ is nonexpansive, then due to Lemma 2.4, the variational inequality (3.3) has a unique solution x' on the set F. We will show that the sequence $\{x_n\}_{n=1}^\infty$ given by (1.8) converges strongly to x'.

STEP 1. The sequences $\{x_n\}_{n=1}^{\infty}$, $\{f(x_n)\}_{n=1}^{\infty}$, $\{T_nx_n\}_{n=1}^{\infty}$, and $\{AT_nx_n\}_{n=1}^{\infty}$, are bounded. Indeed, let $p \in F$ and $S_n := (I - \alpha_n\mu A)T_n$. Using (3.2), we then have

$$\begin{aligned}\|x_{n+1} - p\| &= \|\alpha_n\gamma f(x_n) + S_nx_n - p\| \\ &= \|\alpha_n\gamma f(x_n) - \alpha_n\mu Ap + S_nx_n - S_np\| \\ &\le \alpha_n\gamma\alpha\|x_n - p\| + \alpha_n\|\gamma f(p) - \mu Ap\| + (1 - \alpha_n\tau)\|x_n - p\| \\ &= (1 - \alpha_n(\tau - \gamma\alpha))\|x_n - p\| + \alpha_n(\tau - \gamma\alpha)\frac{\|\gamma f(p) - \mu Ap\|}{\tau - \gamma\alpha}. \qquad (3.4)\end{aligned}$$

Since $0 < \gamma < \frac{\tau}{\alpha}$, $0 < \alpha_n < 1$, and $\lim_{n\to\infty} \alpha_n = 0$ then there exists $n_0 > 0$ such that $0 < \alpha_n(\tau - \gamma\alpha) < 1$ for all $n \ge n_0$. We then obtain

$$\|x_{n+1} - p\| \le \max\left\{\|x_n - p\|, \frac{\|\gamma f(p) - \mu Ap\|}{\tau - \gamma\alpha}\right\}, \quad \forall n \ge n_0. \qquad (3.5)$$

Therefore, we get

$$\|x_{n+1} - p\| \le \max\left\{\|x_{n_0} - p\|, \frac{\|\gamma f(p) - \mu Ap\|}{\tau - \gamma\alpha}\right\}, \quad \forall n \ge n_0, \qquad (3.6)$$

which means, $\{x_n\}_{n=1}^{\infty}$ is a bounded sequence.

From the following inequality

$$\|f(x_n) - f(p)\| \le \alpha\|x_n - p\|,$$

$$\|T_nx_n - p\| = \|T_nx_n - T_np\| \le \|x_n - p\|,$$

$$\|AT_nx_n - Ap\| = \|AT_nx_n - AT_np\| \le k\|T_nx_n - T_np\| \le k\|x_n - p\|,$$

it follows that the sequences $\{f(x_n)\}_{n=1}^{\infty}$, $\{T_nx_n\}_{n=1}^{\infty}$, $\{AT_nx_n\}_{n=1}^{\infty}$, are bounded.

STEP 2. One has $\lim_{n\to\infty} \|x_{n+1} - T_nx_n\| = 0$. Indeed, from $\lim_{n\to\infty} \alpha_n = 0$, (1.8), and Step 1 it follows that

$$\lim_{n\to\infty} \|x_{n+1} - T_nx_n\| = \lim_{n\to\infty} \alpha_n\|\gamma f(x_n) - \mu AT_nx_n\| = 0$$

STEP 3. For the sequence $\{x_n\}$ we have $\lim_{n\to\infty} \|x_{n+N} - x_n\| = 0$. Indeed, noting that $T_{n+N-1} = T_{n-1}$ and

$$S_{n+N-1}x_{n-1} - S_{n-1}x_{n-1} = \mu(\alpha_{n+N-1} - \alpha_{n-1})AT_{n-1}x_{n-1}$$

we then obtain

$$\begin{aligned}\|x_{n+N} - x_n\| &\le \|\alpha_{n+N-1}\gamma f(x_{n+N-1}) - \alpha_{n-1}\gamma f(x_{n-1})\| \\ &\quad +\|S_{n+N-1}x_{n+N-1} - S_{n-1}x_{n-1}\| \\ &\le \alpha_{n+N-1}\gamma\alpha\|x_{n+N-1} - x_{n-1}\| + \gamma|\alpha_{n+N-1} - \alpha_{n-1}|\|f(x_{n-1})\| \\ &\quad +\|S_{n+N-1}x_{n+N-1} - S_{n+N-1}x_{n-1}\| \\ &\quad +\mu|\alpha_{n+N-1} - \alpha_{n-1}|\|AT_{n-1}x_{n-1}\| \\ &\le (1 - \alpha_{n+N-1}(\tau - \gamma\alpha))\|x_{n+N-1} - x_{n-1}\| \\ &\quad +|\alpha_{n+N-1} - \alpha_{n-1}|\Big(\gamma\|f(x_{n-1})\| + \mu\|AT_{n-1}x_{n-1}\|\Big) \\ &= (1 - \alpha_{n+N-1}(\tau - \gamma\alpha))\|x_{n+N-1} - x_{n-1}\| \\ &\quad +\alpha_{n+N-1}(\tau - \gamma\alpha)\beta_{n+N-1},\end{aligned}$$

where

$$\beta_{n-N+1} := \frac{|\alpha_{n+N-1} - \alpha_{n-1}|}{\alpha_{n+N-1}} \cdot \frac{\gamma\|f(x_{n-1})\| + \mu\|AT_{n-1}x_{n-1}\|}{\tau - \gamma\alpha}.$$

Letting $a_{n+N} := \|x_{n+N} - x_n\|$, one then gets

$$a_{n+N} \leq (1 - \alpha_{n+N-1}(\tau - \gamma\alpha))a_{n+N-1} + \alpha_{n+N-1}(\tau - \gamma\alpha)\beta_{n+N-1}, \tag{3.7}$$

where

$$\limsup_{n\to\infty} \beta_n = \limsup_{n\to\infty} \frac{|\alpha_{n+N-1} - \alpha_{n-1}|}{\alpha_{n+N-1}} \cdot \frac{\gamma\|f(x_{n-1})\| + \mu\|AT_{n-1}x_{n-1}\|}{\tau - \gamma\alpha} = 0$$

According to Lemma 2.1, we obtain

$$\lim_{n\to\infty} a_{n+N} = \lim_{n\to\infty} \|x_{n+N} - x_n\| = 0.$$

STEP 4. We have $\lim\limits_{n\to\infty} \|x_n - T_{n+N-1}T_{n+N-2}\cdots T_n x_n\| = 0$. Indeed, noting that T_n is a nonexpansive mapping and using Step 2 one has the following

$$\begin{array}{rcl}
\|x_{n+N} - T_{n+N-1}x_{n+N-1}\| & \to & 0, \\
\|T_{n+N-1}x_{n+N-1} - T_{n+N-1}T_{n+N-2}x_{n+N-2}\| & \to & 0, \\
\|T_{n+N-1}T_{n+N-2}x_{n+N-2} - T_{n+N-1}T_{n+N-2}T_{n+N-3}x_{n+N-3}\| & \to & 0, \\
 & \vdots & \\
\|T_{n+N-1}T_{n+N-2}\cdots T_{n+1}x_{n+1} - T_{n+N-1}T_{n+N-2}T_{n+N-3}\cdots T_n x_n\| & \to & 0,
\end{array}$$

as $n \to \infty$. Hence, we obtain

$$\|x_{n+N} - T_{n+N-1}T_{n+N-2}T_{n+N-3}\cdots T_n x_n\| \to 0, \quad n \to \infty.$$

Since $\|x_{n+N} - x_n\| \to 0$ as $n \to \infty$ (see Step 3) we then get

$$\lim_{n\to\infty} \|x_n - T_{n+N-1}T_{n+N-2}\cdots T_n x_n\| = 0.$$

STEP 5. We want to show that $\limsup\limits_{n\to\infty}\Big\langle(\gamma f - \mu A)x', x_n - x'\Big\rangle \leq 0$, where x' is a unique solution of the variational inequality (3.3) for the nonexpansive mapping $T_N T_{N-1}\cdots T_1 : H \to H$ on the set F.

Let $\{x_{n_k}\}_{k=1}^{\infty}$ be a subsequence of $\{x_n\}_{n=1}^{\infty}$ such that

$$\limsup_{n\to\infty}\Big\langle(\gamma f - \mu A)x', x_n - x'\Big\rangle = \lim_{k\to\infty}\Big\langle(\gamma f - \mu A)x', x_{n_k} - x'\Big\rangle.$$

Since $\{x_{n_k}\}_{k=1}^{\infty}$ is bounded and H is a real Hilbert space, then there exists a subsequence $\{x_{n_{k_m}}\}_{m=1}^{\infty}$ of $\{x_{n_k}\}_{k=1}^{\infty}$ such that $\{x_{n_{k_m}}\}_{m=1}^{\infty}$ converges weakly to some point $y \in H$. Without loss any generality, we may assume that n_{k_m} are such kind of numbers that $T_{n_{k_m}} = T_{i_0}$ for some $i_0 \in \{1, 2, \cdots, N\}$ and for all $m \in \mathbb{N}$. Then, from Step 4, it follows that

$$\lim_{n\to\infty} \|x_{n_{k_m}} - T_{i_0+N-1}T_{i_0+N-2}\cdots T_{i_0}x_{n_{k_m}}\| = 0.$$

So due to Lemma 2.3, we have $y \in F(T_{i_0+N-1}T_{i_0+N-2}\cdots T_{i_0}) = F$. Therefore,

$$\begin{array}{rcl}
\limsup\limits_{n\to\infty}\Big\langle(\gamma f - \mu A)x', x_n - x'\Big\rangle & = & \lim\limits_{m\to\infty}\Big\langle(\gamma f - \mu A)x', x_{n_{k_m}} - x'\Big\rangle \\
 & = & \Big\langle(\gamma f - \mu A)x', y - x'\Big\rangle \leq 0.
\end{array}$$

Step 6. The sequence $\{x_n\}$ converges to x', i.e. $\lim_{n\to\infty}\|x_n - x'\| = 0$. Indeed, using Lemma 2.2 and (3.4) we obtain

$$\begin{aligned}
\|x_{n+1}-x'\|^2 &= \|S_n x_n - S_n x' + \alpha_n(\gamma f(x_n) - \mu ATx')\|^2 \\
&\le \|S_n x_n - S_n x'\|^2 + 2\alpha_n\Big\langle \gamma f(x_n) - \mu A x', x_{n+1}-x'\Big\rangle \\
&\le (1-\alpha_n\tau)^2\|x_n - x^{'}\|^2 + 2\alpha_n\gamma\Big\langle f(x_n) - f(x'), x_{n+1}-x'\Big\rangle \\
&\quad +2\alpha_n\Big\langle (\gamma f(x_n) - \mu A)x', x_{n+1}-x'\Big\rangle \\
&\le (1-\alpha_n\tau)^2\|x_n - x^{'}\|^2 + 2\alpha_n\gamma\alpha\|x_n - x'\|\|x_{n+1}-x'\| \\
&\quad +2\alpha_n\Big\langle (\gamma f(x_n) - \mu A)x', x_{n+1}-x'\Big\rangle \\
&\le (1-\alpha_n\tau)^2\|x_n - x^{'}\|^2 + 2\alpha_n(1-\alpha_n(\tau-\gamma\alpha))\gamma\alpha\|x_n - x'\|^2 \\
&\quad +2\alpha_n^2\gamma\alpha\|\gamma f(x') - \mu A x'\| + 2\alpha_n\Big\langle (\gamma f(x_n) - \mu A)x', x_{n+1}-x'\Big\rangle \\
&\le (1-2\alpha_n(\tau-\gamma\alpha))\|x_n - x^{'}\|^2 + 2\alpha_n(\tau-\gamma\alpha)\beta_n,
\end{aligned}$$

where

$$\begin{aligned}
\beta_n &= \frac{\Big\langle (\gamma f(x_n) - \mu A)x', x_{n+1}-x'\Big\rangle}{\tau-\gamma\alpha} \\
&\quad +\alpha_n\left[\left(\frac{\tau^2}{2(\tau-\gamma\alpha)} - \gamma\alpha\right)\|x_n - x'\|^2 + \frac{\gamma\alpha}{\tau-\gamma\alpha}\|\gamma f(x') - \mu A x'\|\right].
\end{aligned}$$

Letting $a_n := \|x_n - x'\|^2$, we then have

$$a_{n+1} \le (1-2\alpha_n(\tau-\gamma\alpha))a_n + 2\alpha_n(\tau-\gamma\alpha)\beta_n. \tag{3.8}$$

Since $\limsup_{n\to\infty}\Big\langle (\gamma f - \mu A)x', x_{n+1}-x'\Big\rangle \le 0$ (see Step 5) one can get

$$\limsup_{n\to\infty}\beta_n \le 0.$$

Therefore, according to Lemma 2.1, we obtain that

$$\lim_{n\to\infty} a_n = \lim_{n\to\infty}\|x_n - x'\|^2 = 0,$$

which implies that $\lim_{n\to\infty}\|x_n - x'\| = 0$. This completes the proof. □

Note that if $f(x) = \gamma u$ for some $\gamma \in (0,1)$ and $u \in H$ we recover Xu's [26] result. Our results also generalize the corresponding results of Marino and Xu [10], Tian [24], Yamada [27].

ACKNOWLEDGEMENT

This work was done while the first named author (F.M.) was visiting the Abdus Salam International Centre for Theoretical Physics, Trieste, Italy as a Junior Associate. He would like to thank the Centre for hospitality and financial support.

References

[1] Bauschke H. H. The Approximation of fixed points of compositions of nonexpansive mappings in Hilbert spaces, *J. Math. Anal. Appl.* 202 (1996) 150-159.

[2] Byrne C. A unified treatment of some iterative algorithms in signal processing and image construction, *Inverse problems* 20 (2004) 103-120.

[3] Chidume C. E. and Ofoedu E. U. A new iteration process for finite families of generalized Lipschitz pseudo-contractive and generalized Lipschitz accretive mappings, *Nonlinear Analysis; TMA.*, In press, accepted manuscript, available online 27 June 2007.

[4] Chidume C. E.and Ofoedu E. U. Approximation of common fixed points for finite families of total asymptotically nonexpansive mappings, *J. Math. Anal. Appl.*, 333(1) (2007), 128-141.

[5] Goebel K. Kirk W. A. *Topics in metric fixed point theory*, Cambridge Studies in Advanced Mathematics 28, CUP, Cambridge, New York, Port Chester, Melbourne, Sydney, 1990.

[6] Halpern B. Fixed points of nonexpanding maps, *Bulletin of the American Mathematical Society,* 73 (1967), 957-961.

[7] Ishikawa S. Fixed point by a new iteration method, *Proc. Amer. Math. Soc.* 44 (1974), 147-150.

[8] Lions P. L. Approximation de points fixes de contractions, Computes rendus de lacademie des sciences, serie I-mathematique, 284 (1997) 1357-1359.

[9] Mann W. R. Mean value methods in iteration, *Proc. Amer. Math. Soc.* 4 (1953) 506-510.

[10] Marino G. and Xu H. K. A general iterative method for nonexpansive mappings in Hilbert spaces, *J. Math. anal. Appl.* 318 (2006) 43-52.

[11] Megginson R. E. An introduction to Banach space theory, Springer-Verlag, New York, Inc., 1998.

[12] Moudafi A. Viscosity approximation metheds for fixed-points problems, *J. Math. Anal. Appl.* 241 (2000) 46-55.

[13] Ofoedu E. U. Iterative approximation of a common zero of a countably infinite family of m-accretive operators in Banach spaces, *Hindawi-Fixed point theory and applications,* doi:10.1155/2008/325792.

[14] Ofoedu E. U. and Shehu Y. Iterative construction of a common fixed point of finite families of nonlinear mappings, in press, to appear in Vol. 16 No 1 of *Journal of Applied Analysis* (2010).

[15] Ofoedu E. U., Shehu Y., and Ezeora J. N. Solution by iteration of nonlinear variational inequalities involving finite family of asymptotically nonexpansive mappings and monotone mappings, *PanAmer. Math. J.* 18 (2008) (4), 61-75.

[16] Polilchuk C. I. and Mammone R. J. Image recovery by convex projections using a least-square constraint, *J. Opt. Soc. Amer.* A7 (1990) 517-521.

[17] Reich S. Weak convergence theorems for nonexpansive mappings in Banach spaces, *J. Math. Anal. Appl.* 67 (1979), 274-276.

[18] Reich S. Extension problems for accretive sets in Banach spaces, *J. Functional Analysis* 26 (1977), 378-395.

[19] S. Reich, Strong convergence theorems for resolvents of accretive operators in Banach spaces, *J. Math. Anal. Appl.*, 75 (1980), 287-292.

[20] Rhoades B. E. Fixed point iterations for certain nonlinear mappings, *J Math. Anal. Appl.* 183 (1994) 118-120.

[21] Rockefellar R. T. Monotone operators and proximal point algorithm, SIAM, *J. Control Optim.* 14 (1976), 877-898.

[22] Schu J. Iterative construction of fixed points of asymptotically nonexpansive mappings, *J. Math. Anal. Appl.* 158 (1991), 407-413.

[23] Schu J. Weak and strong convergence of fiexd points of asymptotically nonexpansive mappings, *Bull. Austral. Math. Soc.* 43 (1991), 153-159.

[24] Tian M. A general iterative algorithm for nonexpansive mappings in Hilbert spaces, *Nonlinear Analysis* (2010), doi: 10.1016/j.na.2010.03.058.

[25] Wittmann R. Approximation of fixed points of nonexpansive mappings, *Archiv der mathematik,* 58 (1992) 486-491.

[26] Xu H. K. An iterative approach to quadratic optimization, *Journal of Optimization Theory and Applications*, 116 (2003) no.3, 659-678.

[27] Yamada I. The hybrid steepest decent for the variational inequality problems over the itersection of fixed points sets of nonexpansive mapping: in D. Butnariu, Y. Censor, S. Reich (Eds), Inherently parallel algorithme in feasibility and optimization and their application, Elsevier, New York, (2001) 473-504.

[28] Youla D. On deterministic convergence of iterations of related projection operators, *J. Vis. Commun.* Image Represent 1 (1990), 12-20.

Farrukh Mukhamedov, Department of Computational & Theoretical Sciences, Faculty of Sciences, International Islamic University Malaysia, P.O. Box, 141, 25710, Kuantan, Pahang, Malaysia
E-mail address: `far75m@yandex.ru`

Mansoor Saburov, Department of Computational & Theoretical Sciences, Faculty of Science, International Islamic University Malaysia, P.O. Box, 141, 25710, Kuantan, Pahang, Malaysia
E-mail address: `msaburov@gmail.com`